

## Performance




Table 2\*

### Effect of Supplementation Strategy on Feed Intake and Growth Performance

#### Low Protein Forage Study

	Control	FIAD 12	FIAD Range 30 MF
<b>Prairie Hay (5% Protein)</b>			
Daily Gain, lbs	-0.57	-0.39	-0.22
Hay Intake, lbs	12.1	11.5	13.4
Block Intake, lbs	-	0.90	1.05
<b>Prairie Hay with 5 lbs/day Alfalfa (8% Protein)</b>			
Daily Gain, lbs	0.81	0.89	1.14
Hay Intake, lbs	18.4	18.0	18.8
Block Intake, lbs	-	0.72	0.80

\* Kansas State University



Protein frequently is the most limiting nutrient for cattle wintered on poor to moderate quality harvested forages as well as for mature cows and growing cattle grazing dormant grass pastures and crop residues. Ample amounts of protein are required in order for cattle to efficiently harvest energy from forage-based feeds. Protein is needed to support a healthy, active population of fiber-digesting microorganisms in the stomach. Consequently, when dietary protein is limiting, fiber digestion is not optimal. Poor digestion results in inefficient energy use, and often decreases feed intake too. The combined effect of these factors is a significant decrease in overall energy balance for the animal. This can limit growth rate of young animals or lead to poor body condition scores in reproducing cows and heifers.

### Cattle Performance Trial

A study was performed at the Kansas State University Beef Cattle Research Center using 175 crossbred beef heifers averaging 683 pounds. Heifers were fed prairie hay free-choice, or the same hay supplemented with 5 pounds daily of alfalfa hay. Heifers within each forage group had access to The Feed In A Drum blocks or no supplement. Heifers were fed the forage-based diets for 89 days, and had free access to white salt and water at all times.

Daily gains, feed intakes, and efficiencies of heifers are shown in Table 2. Heifers fed prairie hay with no alfalfa actually lost weight, indicating that the hay had limited energy content. Performance was improved substantially by feeding blocks. In particular, the higher protein content of Range 30 MF improved forage intake and reduced weight loss compared to the unsupplemented controls.

Feeding five pounds of alfalfa daily effectively increased overall energy intake by heifers, allowing for a substantial improvement in gain compared to heifers fed only the prairie hay. Feeding The Feed In A Drum blocks resulted in additional boosts in gain and efficiency, illustrating that The Feed In A Drum supplements are effective over a range of forage qualities.